

# The burden and epidemiology of community-acquired central nervous system infections: a multinational study

H. Erdem<sup>1,2</sup> · A. Inan<sup>3</sup> · E. Guven<sup>4</sup> · S. Hargreaves<sup>5</sup> · L. Larsen<sup>6</sup> · G. Shehata<sup>7</sup> · E. Pernicova<sup>8,9</sup> · E. Khan<sup>10</sup> · L. Bastakova<sup>11</sup> · S. Namani<sup>12</sup> · A. Harxhi<sup>13</sup> · T. Roganovic<sup>14</sup> · B. Lakatos<sup>15</sup> · S. Uysal<sup>16</sup> · O. R. Sipahi<sup>17</sup> · A. Crisan<sup>18</sup> · E. Miftode<sup>19</sup> · R. Stebel<sup>11</sup> · B. Jegorovic<sup>20</sup> · Z. Fehér<sup>21</sup> · C. Jekkel<sup>15</sup> · N. Pandak<sup>22</sup> · A. Moravveji<sup>23</sup> · H. Yilmaz<sup>24</sup> · A. Khalifa<sup>25</sup> · U. Musabak<sup>26</sup> · S. Yilmaz<sup>27</sup> · A. Jouhar<sup>25</sup> · N. Oztoprak<sup>28</sup> · X. Argemi<sup>29</sup> · M. Baldeyrou<sup>29</sup> · G. Bellaud<sup>30</sup> · R. V. Moroti<sup>31</sup> · R. Hasbun<sup>32</sup> · L. Salazar<sup>32</sup> · R. Tekin<sup>33</sup> · A. Canestri<sup>30</sup> · L. Čalkić<sup>34</sup> · L. Praticò<sup>35</sup> · F. Yilmaz-Karadag<sup>36</sup> · L. Santos<sup>37</sup> · A. Pinto<sup>37</sup> · F. Kaptan<sup>38</sup> · P. Bossi<sup>39</sup> · J. Aron<sup>39</sup> · A. Duissenova<sup>40</sup> · G. Shopayeva<sup>40</sup> · B. Utaganov<sup>40</sup> · S. Grgic<sup>41</sup> · G. Ersoz<sup>42</sup> · A. K. L. Wu<sup>43</sup> · K. C. Lung<sup>43</sup> · A. Bruzsa<sup>15</sup> · L. B. Radic<sup>44</sup> · H. Kahraman<sup>17</sup> · M. Momen-Heravi<sup>45</sup> · S. Kulzhanova<sup>46</sup> · F. Rigo<sup>47</sup> · M. Konkayeva<sup>46</sup> · Z. Smagulova<sup>46</sup> · T. Tang<sup>48</sup> · P. Chan<sup>49</sup> · S. Ahmetagic<sup>50</sup> · H. Porobic-Jahic<sup>50</sup> · F. Moradi<sup>51</sup> · S. Kaya<sup>52</sup> · Y. Cag<sup>53</sup> · A. Bohr<sup>54</sup> · C. Artuk<sup>55</sup> · I. Celik<sup>56</sup> · M. Amsilli<sup>57</sup> · H. C. Gul<sup>55</sup> · A. Cascio<sup>58</sup> · M. Lanzafame<sup>47</sup> · M. Nassar<sup>59</sup>

Received: 12 February 2017 / Accepted: 22 March 2017 / Published online: 10 April 2017  
© Springer-Verlag Berlin Heidelberg 2017

**Abstract** Risk assessment of central nervous system (CNS) infection patients is of key importance in predicting likely pathogens. However, data are lacking on the epidemiology globally. We performed a multicenter study to understand the burden of community-acquired CNS (CA-CNS) infections

between 2012 and 2014. A total of 2583 patients with CA-CNS infections were included from 37 referral centers in 20 countries. Of these, 477 (18.5%) patients survived with sequelae and 227 (8.8%) died, and 1879 (72.7%) patients were discharged with complete cure. The most frequent infecting

**Electronic supplementary material** The online version of this article (doi:10.1007/s10096-017-2973-0) contains supplementary material, which is available to authorized users.

✉ H. Erdem  
hakanerdem1969@yahoo.com

- 1 Principal Coordinator of ID-IRI, Ankara, Turkey
- 2 Department of Infectious Diseases and Clinical Microbiology, Gulhane Medical Academy, 06010 Etlik, Ankara, Turkey
- 3 Department of Infectious Diseases and Clinical Microbiology, Haydarpasa Numune Training and Research Hospital, Istanbul, Turkey
- 4 Beytepe Murat Erdi Eker State Hospital, Ankara, Turkey
- 5 International Health Unit, Section of Infectious Diseases and Immunity, Commonwealth Building, Hammersmith Campus, Imperial College London, London, UK
- 6 Department of Infectious Diseases Q, Odense University Hospital, Odense, Denmark

- 7 Department of Neurology and Psychiatry, Assiut University Hospital, Assiut, Egypt
- 8 Avenier, Centres for Vaccination and Travel Medicine, Prague, Czech Republic
- 9 Faculty Hospital Brno, Department of Infectious Diseases, Brno, Czech Republic
- 10 Shifa International Hospital, Islamabad, Pakistan
- 11 Faculty Hospital Brno, Department of Infectious Diseases and Masaryk University Faculty of Medicine, Brno, Czech Republic
- 12 Infectious Diseases Clinic, University Clinical Center of Kosovo, Prishtina, Kosovo
- 13 Service of Infectious Disease, University Hospital Center of Tirana, Tirana, Albania
- 14 Infectious Diseases Clinic, University Hospital Clinical Center Banja Luka, Banja Luka, Bosnia and Herzegovina

pathogens in this study were *Streptococcus pneumoniae* ( $n = 206$ , 8%) and *Mycobacterium tuberculosis* ( $n = 152$ , 5.9%). Varicella zoster virus and *Listeria* were other common pathogens in the elderly. Although staphylococci and *Listeria* resulted in frequent infections in immunocompromised patients, cryptococci were leading pathogens in human

immunodeficiency virus (HIV)-positive individuals. Among the patients with any proven etiology, 96 (8.9%) patients presented with clinical features of a chronic CNS disease. Neurosyphilis, neurobrucellosis, neuroborreliosis, and CNS tuberculosis had a predilection to present chronic courses. *Listeria monocytogenes*, *Staphylococcus aureus*,

- 15 Department of Infectious Diseases, Saint Laszlo Hospital, Budapest, Hungary
- 16 Department of Infectious Diseases and Clinical Microbiology, Seyfi Demirsoy State Hospital, Buca, Izmir, Turkey
- 17 Department of Infectious Diseases and Clinical Microbiology, Ege University School of Medicine, Izmir, Turkey
- 18 Department of Infectious Diseases, Victor Babes University of Medicine and Pharmacy, Timisoara, Romania
- 19 Hospital of Infectious Diseases, Gr. T. Popa University of Medicine and Pharmacy, Iasi, Romania
- 20 Clinic for Infectious and Tropical Diseases, Clinical Centre of Serbia, Faculty of Medicine, University of Belgrade, Belgrade, Serbia
- 21 Department of Infectious Diseases, Markusovszky University Teaching Hospital, Szombathely, Hungary
- 22 General Hospital Slavonski Brod, Department for Infectious Diseases, School of Medicine, University of Split, Split, Croatia
- 23 Social Determinants of Health Research Center, Department of Community Medicine, Kashan University of Medical Sciences, Kashan, Iran
- 24 Department of Infectious Diseases and Clinical Microbiology, Ondokuz Mayıs University School of Medicine, Samsun, Turkey
- 25 Department of Neurology, Damascus Hospital, Damascus, Syria
- 26 Department of Immunology and Allergy, Losante Hospital, Ankara, Turkey
- 27 Gulhane Medical Academy, Blood Bank, Clinical Microbiology Division, Ankara, Turkey
- 28 Antalya Education and Research Hospital, Antalya, Turkey
- 29 Infectious Diseases Department, Nouvel Hôpital Civil, Strasbourg, France
- 30 Department of Infectious Diseases, Tenon University Hospital, Paris, France
- 31 Carol Davila University of Medicine and Pharmacy and Matei Bals National Institute for Infectious Diseases, Bucharest, Romania
- 32 Medical School, Department of Infectious Diseases, The University of Texas Health Science Center at Houston, Houston, TX, USA
- 33 Department of Infectious Diseases and Clinical Microbiology, Dicle University School of Medicine, Diyarbakir, Turkey
- 34 Department of Infectious Diseases, Zenica Cantonal Hospital, Zenica, Bosnia and Herzegovina
- 35 University Division of Infectious and Tropical Diseases, Piazza Spedali Civili, 25123 Brescia, Italy
- 36 Goztepe Training and Research Hospital, Department of Infectious Diseases and Clinical Microbiology, Medeniyet University, Istanbul, Turkey
- 37 Infectious Diseases Service, Centro Hospitalar São João and Faculty of Medicine, University of Porto, Porto, Portugal
- 38 Department of Infectious Diseases and Clinical Microbiology, Katip Celebi University School of Medicine, Izmir, Turkey
- 39 Department Maladies Infectieuses, Institut Pasteur de Paris—HPA, Paris, France
- 40 Department of Infectious and Tropical Diseases, Asfendiyarov Kazakh National Medical University, Almaty, Kazakhstan
- 41 Clinic for Infectious Diseases, University Hospital of Mostar, Mostar, Bosnia and Herzegovina
- 42 Department of Infectious Diseases and Clinical Microbiology, Mersin University School of Medicine, Mersin, Turkey
- 43 Department of Clinical Pathology, Pamela Youde Nethersole Eastern Hospital, Hong Kong SAR, China
- 44 Department of Infectious Diseases, General Hospital Dubrovnik, Dubrovnik, Croatia
- 45 Department of Infectious Diseases, Social Determinants of Health Research Center, Kashan University of Medical Sciences, Kashan, Iran
- 46 Department of Infectious Diseases, Astana Medical University, Astana, Kazakhstan
- 47 Unit of Infectious Diseases, Azienda Ospedaliera Universitaria Integrata, Verona, Italy
- 48 Infectious Diseases Team, Department of Medicine, Queen Elizabeth Hospital, Hong Kong, China
- 49 Neurology Team, Department of Medicine, Queen Elizabeth Hospital, Hong Kong, China
- 50 University Clinical Center Tuzla, Tuzla, Bosnia and Herzegovina
- 51 Infectious and Tropical Diseases Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran
- 52 Department of Infectious Diseases and Clinical Microbiology, Karadeniz Technical University School of Medicine, Trabzon, Turkey
- 53 School of Medicine, Department of Infectious Diseases and Clinical Microbiology, Medeniyet University, Istanbul, Turkey
- 54 Institute of Inflammation Research, Department of Infectious Diseases and Rheumatology, Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark
- 55 Department of Infectious Diseases and Clinical Microbiology, Gulhane Training and Research Hospital, Ankara, Turkey
- 56 Department of Infectious Diseases and Clinical Microbiology, Kayseri Training and Research Hospital, Kayseri, Turkey
- 57 Infectious Diseases Unit, CHU Bicêtre, Paris, France
- 58 Department of Health Promotion Sciences and Mother and Child Care “G. D’Alessandro”, University of Palermo, Palermo, Italy
- 59 Infection Control Department, Saudi German Hospital Group, Jeddah, Saudi Arabia

*M. tuberculosis*, and *S. pneumoniae* were the most fatal forms, while sequelae were significantly higher for herpes simplex virus type 1 ( $p < 0.05$  for all). Tackling the high burden of CNS infections globally can only be achieved with effective pneumococcal immunization and strategies to eliminate tuberculosis, and more must be done to improve diagnostic capacity.

## Introduction

The central nervous system (CNS) can be infected with numerous infectious agents, which result in significant morbidity and mortality, despite the effectiveness of available antimicrobials [1–5]. In large international cohorts, the case–fatality rates for adults with pneumococcal, tuberculous, and herpetic CNS infections are 14, 17, and 10%, respectively, despite treatment [6–8]. Thus, risk assessment in this subgroup of patients is of key importance in predicting the likely pathogen leading to rational treatment. Although there are considerable data on the epidemiology of these CNS infectious syndromes, data are lacking to facilitate our understanding of the cumulative epidemiology of community-acquired CNS (CA-CNS) infections. We, therefore, performed a multicenter study to provide an insight into CA-CNS infections and to better understand the burden of infection in community settings, and to explore risk assessment data for the management of these infections.

## Materials and methods

### Study design

We designed a multicenter, retrospective, cross-sectional study performed in the infectious diseases and neurology departments of participating referral hospitals. The study was led by the Infectious Diseases International Research Initiative (ID-IRI). We approached >50 survey sites in >25 countries to participate in this study. Data were collected retrospectively during the period from 1st May 2012 to 1st May 2014 and included all CA-CNS infection patients over the age of 15 years hospitalized at the survey sites. All epidemiological, clinical, laboratory, therapeutic, and outcome data were collected using a standardized questionnaire (Excel 2010). The participant centers submitted their individual datasets via email to the lead hospital for analysis. Patient confidentiality was considered and the study was approved ethically by the institutional review board of Fatih Sultan Mehmet Education and Research Hospital in Istanbul.

## Definitions

The diagnoses were made according to the clinical suspicion of the examining clinicians based on compatible clinical presentations of the patients with CA-CNS infections, and in accordance with standard definitions [2–5, 9]. The CA-CNS infections were defined as infections affecting the CNS, and were acquired in daily life. These infectious disorders were not associated to infections in the healthcare settings.

Chronic CA-CNS infection	Patients with symptoms for more than four weeks at the time of diagnosis were included in this category [10].
Elderly classification	Patients over the age of 64 years were classified as elderly [11].
Primary immunosuppression	Congenital immunodeficiency diseases are classified in this category [12].
Secondary immunosuppression	Immunodeficiency conditions depending on underlying immune diseases or due to immunosuppressive treatments are defined as secondary [12].
Probable immunosuppression	Patients with probable immunosuppressive conditions like human immunodeficiency virus (HIV; immune status unknown), diabetes mellitus, malignancy, pregnancy, and others were included in this group. Patients were included in this group when they did not have detailed data in our database in determining the level of immunosuppression.
Adverse clinical outcome	Either death or sequelae were defined as adverse clinical outcomes.
Data sources and measurements	The microbiological diagnoses of patients compatible with CNS infections were made according to the microbiological techniques below and are presented in detail in Table 1:
	<ul style="list-style-type: none"> <li>a) Neurosyphilis: CSF VDRL, serum VDRL/RPR, fluorescent treponemal antibody absorption (FTA-ABS), <i>T. pallidum</i> particle agglutination (TPPA)/<i>T. pallidum</i> hemagglutination test (TPHA), <i>T. pallidum</i> microhemagglutination (MHA-TP) [13].</li> <li>b) Neuroborreliosis: Molecular testing and serology [14].</li> <li>c) Neurobrucellosis: Non-specific CSF culture, blood culture, CSF Rose Bengal test, CSF</li> </ul>

**Table 1** The efficacy of the microbiological methods used in the diagnosis of community-acquired central nervous system (CA-CNS) infections

Bacteria/fungus	Tested	Positive	Positivity (%)	Overall (%)	Bacteria/fungus	Tested	Positive	Positivity (%)	Overall (%)
<i>T. pallidum</i>					<i>Fusobacterium</i> spp.				
CSF VDRL	179	17	9.49	0.65	Culture (CSF)	1414	1	0.07	0.04
Serum VDRL/RPR	346	6	1.73	0.2	Molecular	222	1	0.45	0.04
CSF FTA-ABS	10	1	10	0.04	Subtotal cases		2		0.08
Serum FTA-ABS	35	5	14.28	0.19	<i>Borrelia</i> spp.				
Subtotal cases		24		0.93	Molecular	128	13	10.1	0.5
<i>Brucella</i> spp.					Serology	508	25	4.92	0.97
Culture (CSF)	1414	3	0.21	0.1	Subtotal cases		38		1.47
Culture (blood)	1565	1	0.06	0.04	<i>Enterococcus</i> spp.				
CSF Rose Bengal	90	6	6.67	0.2	Culture (CSF)	1414	4	0.28	0.15
CSF Wright aggl. test	81	8	9.87	0.3	Culture (blood)	1565	2	0.13	0.07
Serum Rose Bengal	188	11	5.85	0.4	Subtotal cases		5		0.19
Serum Wright aggl. test	207	13	6.28	0.5	<i>Micrococcus</i> spp. (blood)				
CSF molecular	1	0	0	0	1565	1	0.06		0.04
Serology	508	1	0.19	0.04	<i>Gemella</i> spp. (CSF)				
Subtotal cases		14		0.54	1414	1	0.07		0.04
<i>M. tuberculosis</i>					<i>Peptococcus</i> spp.				
Culture (CSF)	528	38	7.19	1.5	Culture (CSF)	1414	1	0.07	0.04
ARB	566	33	5.83	1.3	Culture (blood)	1565	1	0.06	0.04
PCR	57	10	17.54	0.4	Subtotal cases		1		0.04
Anti-TB treatment		107		4	<i>Peptostreptococcus</i> spp.				
Subtotal cases		152		5.88	Culture (CSF)	1414	2	0.14	0.08
<i>S. pneumoniae</i>					Culture (blood)	1565	1	0.06	0.04
Culture (CSF)	1414	148	10.46	5.7	Subtotal cases		3		0.12
Culture (blood)	1565	85	5.43	3.3	<i>Lactobacillus</i> spp. (CSF)				
Molecular	222	31	13.96	1.2	1414	1	0.07		0.04
Serology	300	40	13.3	1.5	<i>Corynebacteria</i> (CSF)				
Subtotal cases		206		7.97	1414	5	0.35		0.19
<i>Streptococcus</i> spp. (other)					<i>Actinomyces</i> spp. (CSF)				
Culture (CSF)	1414	36	2.54	1.4	1414	5	0.35		0.19
Culture (blood)	1565	20	1.27	0.8	<i>Escherichia coli</i>				
Molecular	222	16	7.20	0.6	1414	6	0.42		0.23
Subtotal cases		55		2.13	Culture (blood)	1565	5	0.32	0.19
<i>Staphylococcus</i> spp.					Subtotal cases		9		0.35
Culture (CSF)	1414	27	1.90	1	<i>Enterobacter</i> spp.				
Culture (blood)	1565	47	3	1.8	Culture (CSF)	1414	2	0.14	0.08
Molecular	222	3	1.35	0.1	Culture (blood)	1565	1	0.06	0.04
Subtotal cases		71		2.75	Subtotal cases		3		0.12
<i>Listeria</i> spp.					<i>Proteus</i> spp. (CSF)				
Culture (CSF)	1414	22	1.55	0.9	1414	2	0.14		0.08
Culture (blood)	1565	12	0.76	0.46	<i>Morganella</i> spp. (CSF)				
Molecular	222	3	1.35	0.1	1414	3	0.21		0.12
Subtotal cases		27		1.04	<i>Campylobacter</i> spp. (CSF)				
<i>Nocardia</i> spp.					1414	1	0.07		0.04
Culture (CSF)	1414	1	0.07	0.04	<i>Aggregatibacter</i> spp.				
Culture (blood)	1565	2	0.13	0.08	Culture (CSF)	1414	2	0.14	0.08
Molecular	205	1	0.48	0.04	Culture (blood)	1565	1	0.06	0.04
Subtotal cases		4		0.15	Subtotal cases		3		0.12
<i>Klebsiella</i> spp.					<i>Pseudomonas</i> spp.				
Culture (CSF)	1414	6	0.42	0.2	Culture (CSF)	1414	4	0.28	0.15
Culture (blood)	1565	2	0.13	0.08	Culture (blood)	1565	4	0.25	0.15
Molecular	222	1	0.45	0.04	Subtotal cases		7		0.27
Subtotal cases		7		0.27	<i>Acinetobacter</i> spp.				
<i>Haemophilus</i> spp.					Culture (CSF)	1414	4	0.28	0.15
Culture (CSF)	1414	7	0.49	0.3	Culture (blood)	1565	1	0.06	0.04
Culture (blood)	1565	1	0.06	0.04	Subtotal cases		4		0.15
Molecular	222	2	0.90	0.08	<i>Moraxella</i> spp. (CSF)				
Serology	300	2	0.67	0.08	1414	2	0.14		0.08
Subtotal cases		10		0.39	<i>Sphingomonas</i> spp. (CSF)				
<i>Capnocytophaga</i> spp.					1414	2	0.14		0.08
Culture (CSF)	1414	2	0.14	0.08	<i>Prevotella</i> spp. (CSF)				
Culture (blood)	1565	2	0.13	0.08	1414	2	0.14		0.08
					<i>Bacteroides</i> spp. (blood)				
					1565	1	0.06		0.04
					<i>Saccharomyces cerevisiae</i> (mol.)				
					222	1	0.45		0.04
					<i>Xanthomonas</i> spp. (mol.)				
					222	1	0.45		0.04
					<i>Aerococcus urinae</i> (mol.)				
					222	1	0.45		0.04
					<i>Cryptococcus</i> spp.				
					Culture (fungal)	509	19	3.73	0.74
					Culture (CSF)	1414	8	0.56	0.31
					Culture (blood)	1565	7	0.44	0.27
					CSF antigen	207	25	12.1	0.96
					Molecular	14	2	14.3	0.08
					Subtotal cases		27		1.04

**Table 1** (continued)

Bacteria/fungus	Tested	Positive	Positivity (%)	Overall (%)	Bacteria/fungus	Tested	Positive	Positivity (%)	Overall (%)
Molecular	222	1	0.45	0.04	<i>Candida</i> spp.				
Subtotal cases		3		0.12	Culture (fungal)	509	2	0.39	0.08
<i>Neisseria</i> spp.					Culture (CSF)	1414	1	0.07	0.08
Culture (CSF)	1414	22	1.55	0.9	Culture (blood)	1565	3	0.19	0.01
Culture (blood)	1565	8	0.51	0.3	Serology	508	1	0.19	0.04
Molecular	222	17	7.65	0.7	Subtotal cases		5		0.19
Serology	300	11	3.67	0.4					
Subtotal cases		45		1.74					
Viruses/parasites	Method		Tested		Positive		Positivity (%)		Overall (%)
HSV-1 (CSF)	Molecular		811		46		5.6		1.8
	Serology		508		2		0.3		0.08
HSV-2 (CSF)	Molecular		791		34		4.3		1.3
HSV-1/2 (CSF)	Serology		508		1		0.2		0.04
HSV (serum)	Molecular		4		1		25		0.04
	Subtotal				74				2.8
EBV (CSF)	Molecular		231		16		6.9		0.62
CMV (CSF)	Molecular		261		2		0.8		0.08
CMV (serum)	Molecular		5		2		40		0.08
	Subtotal				5				0.2
VZV (CSF)	Molecular		548		88		16		3.4
	Serology		508		4		0.8		0.15
	Subtotal				91				3.5
SFTV (CSF)	Molecular		4		1		25		0.04
WNV (CSF)	Molecular		78		8		10		0.3
	Serology		508		13		2.6		0.5
	Subtotal				35				1.35
TBEV	Serology		508		92		18		3.6
Parechovirus (CSF)	Molecular		7		1		14		0.04
Mumps virus (CSF)	Molecular		23		6		26		0.23
Adenovirus (CSF)	Molecular		37		1		2.7		0.04
	Serology		508		1		0.2		0.04
	Serum		7		3		42		0.12
	molecular								
	Subtotal				4				0.15
Enterovirus (CSF)	Molecular		347		88		25		3.4
	Serology		508		1		0.2		0.04
	Serum		9		3		33		0.12
	molecular								
	Subtotal				92				3.6
Measles virus	Serology		508		1		0.2		0.04
HHV6 (CSF)	Molecular		92		3		3.3		0.12
HIV (CSF)	Molecular		24		12		50		0.46
JCV (CSF)	Molecular		22		5		23		0.2
RSV (CSF)	Molecular		5		1		20		0.04
PIV (CSF)	Molecular		5		1		20		0.04
	Serology		508		1		0.2		0.04
	Subtotal				2				0.08
<i>Toxoplasma</i> spp.	Molecular		44		4		9		0.15
(CSF)	Serology		508		1		0.2		0.04
	Serum		3		3		100		0.12
	molecular								
	Subtotal				8				0.3

SFTV Sandfly fever Toscana virus; WNV West Nile virus; TBEV tick-borne encephalitis virus; PIV parainfluenza virus

- d) CNS tuberculosis: CSF culture for tuberculosis, acid fast staining of CSF, molecular testing (MTB

- PCR), and response to anti-tuberculosis treatment in probable cases [16].
- e) Other bacterial infections: Non-specific CSF culture, blood culture, and molecular testing [1, 17].



- f) Cryptococcal CNS disease: Fungal CSF culture, non-specific CSF culture, blood culture, CSF cryptococcal antigen testing, and molecular testing [17].
- g) Candidal CNS disease: Fungal CSF culture, non-specific CSF culture, blood culture, and serological testing [18].
- h) Viral CNS infections: Molecular testing and serology [3, 5].
- i) CNS toxoplasmosis: Molecular testing and serology [5].

The detection of pathogens was directly related to the technical capacity of the participating centers and the diagnostic approaches of the examining clinicians. HIV parameters (e.g., CD4+ T-cell counts and HIV-RNA levels) indicating severity of illness were not in our database. Hence, HIV-positives are included in a separate block.

### Statistical methods

Chi-square, Pearson's chi-square, continuity correction of Pearson, and Fisher's exact test were used to analyze the data. We used a cut-off  $p$ -value of  $p < 0.05$  to indicate significance for all tests. Continuous variables were analyzed by using histograms and the Kolmogorov–Smirnov test for normality. After this separation, a Mann–Whitney  $U$ -test was used for variables that were not distributed normally.

### Results

A total of 2583 patients with CA-CNS infections were included from 37 referral centers in Albania, Bosnia and Herzegovina, Croatia, Czech Republic, Denmark, Egypt, France, Hungary, Hong Kong, Iran, Italy, Kazakhstan, Kosovo, Pakistan, Portugal, Romania, Serbia, Syria, Texas/USA, and Turkey. Table 1 highlights the microbiological techniques and their positivity rates used in the diagnosis.

### Infectious syndromes

There were 2603 infectious CNS syndromes identified in 2583 cases. Patients were classified as meningitis ( $n = 1292$ , 50%), encephalitis ( $n = 208$ , 8%), meningoencephalitis ( $n = 888$ , 34.4%), and meningoencephalomyelitis ( $n = 1$ , 0.04%). There were 144 (5.6%) patients with suppurative intracranial infections. These were brain abscesses ( $n = 97$ ), subdural empyemas ( $n = 14$ ), epidural abscesses ( $n = 31$ ), other abscesses ( $n = 2$ ), other syndromes ( $n = 69$ ), and arachnoiditis

( $n = 2$ ). The distribution of infectious CNS syndromes is presented in Table 2.

### Microbiological data

In this study, a CNS pathogen was disclosed in 1079 (41.8%) patients. However, no infecting pathogen was detected in 1504 (58.2%) patients. The diagnoses were established with either direct or indirect methods, or with the combination of different methods, as indicated in Table 1. We identified *Streptococcus pneumoniae* ( $n = 206$ , 8%) and *Mycobacterium tuberculosis* ( $n = 152$ , 5.9%) as the frequent causative agents. The distribution of infecting pathogens according to the ultimate diagnosis and immunosuppressive states are presented in Tables 3 and 4, respectively. Common pathogens in CA-CNS infections are presented in Fig. 1.

In 77 out of 2583 (3%) patients, multiple etiological agents were detected. In 75 cases, two causative agents were identified and in two patients, three agents were recorded. When these patients were categorized according to clinical presentations, there were 43 cases of meningitis, in which 87 different organisms were identified [*S. pneumoniae* ( $n = 20$ ), *T. pallidum* ( $n = 10$ ), enteroviruses ( $n = 9$ ), cryptococci ( $n = 7$ ), EBV ( $n = 5$ ), streptococci other than pneumococci ( $n = 4$ ), *N. meningitidis* ( $n = 3$ ), VZV ( $n = 3$ ), WNV ( $n = 3$ ), coagulase-negative staphylococci ( $n = 2$ ), *Borrelia* spp. ( $n = 2$ ), HHV-6 ( $n = 1$ ), CMV ( $n = 1$ ), *Brucella* spp. ( $n = 1$ ), *Listeria* spp. ( $n = 1$ )]. In 26 patients with meningoencephalitis, 53 agents were recorded as multiple pathogens [TBEV ( $n = 16$ ), *Borrelia* spp. ( $n = 13$ ), VZV ( $n = 10$ ), *M. tuberculosis* ( $n = 3$ ), HSV ( $n = 3$ ), enteroviruses ( $n = 3$ ), EBV ( $n = 2$ ), *S. pneumoniae* ( $n = 1$ ), coagulase-negative staphylococci ( $n = 1$ ), *Cryptococcus* ( $n = 1$ )]. In five patients with encephalitis, ten agents [HSV-1 ( $n = 4$ ), *S. aureus* ( $n = 1$ ), *S. pneumoniae* ( $n = 1$ ), VZV ( $n = 1$ ), WNV ( $n = 1$ ), EBV ( $n = 1$ ), enterovirus ( $n = 1$ )] were identified. Finally, in three patients with brain abscesses, six pathogens were disclosed [streptococci other than pneumococci ( $n = 2$ ), *Neisseria* spp. ( $n = 2$ ), CMV ( $n = 1$ ), EBV ( $n = 1$ )].

### Age distribution

The mean age of the cases was 47.63 (SD  $\pm$  19.8) years. 446 (17.3%) patients were elderly, with a mean age of 74.58 (SD  $\pm$  7.02) years. Among the patients with any identified pathogen, the numbers of patients over 65 years and less than 65 years of age were 250 (23.2%) and 829 (80.9%), respectively. Although *S. pneumoniae* [59/250–147/829 ( $p = 0.039$ )], *Listeria* spp. [13/250–14/829 ( $p = 0.004$ )], and VZV [41/250–50/829 ( $p < 0.001$ )] caused significantly more infections in the elderly, *M. tuberculosis* [23/250–129/829 ( $p = 0.011$ )], enteroviruses [2/250–89/829 ( $p < 0.001$ )], and

**Table 2** The presentation of the patients according to etiological agents

	No.	Meningitis	Encephalitis	Meningoencephalitis	Brain abscess	Subdural empyema	Epidural abscess	Other abscesses	Other syndromes	Arachnoiditis	Chronic infections		Age (years)	p-Value		
											Agents (n, %)	Patients* (%)				
<i>T. pallidum</i>	24	22	1	0	1	0	0	0	0	0	24	100	18	22	2	0.134
<i>Brucella</i> spp.	14	12	0	1	0	0	1	0	0	0	6	42.8	4.5	13	1	0.210
<i>M. tuberculosis</i>	152	84	4	59	1	0	2	0	3	1	37	24.3	28	129	23	<b>0.011</b>
<i>S. pneumoniae</i>	206	136	4	65	1	2	0	0	0	0	0	0	0	147	59	<b>0.039</b>
<i>Streptococcus</i> spp. (other)	55	22	0	14	16	3	4	0	0	0	0	0	0	45	10	0.0462
<i>S. aureus</i>	48	28	1	4	3	2	9	2	1	0	1	2.1	0.7	32	16	0.125
Co-NS	23	11	1	7	3	0	1	0	1	0	2	8.7	1.5	19	4	0.999
<i>Listeria</i> spp.	27	20	1	6	0	0	0	0	0	0	0	0	0	14	13	<b>0.004</b>
<i>Nocardia</i> spp.	4	0	0	0	4	0	0	0	0	0	1	25	0.7	0	4	
<i>Klebsiella</i> spp.	7	4	0	0	3	0	0	0	0	0	1	14.3	0.7	7	0	
<i>Haemophilus</i> spp.	10	8	0	1	1	0	0	0	0	0	0	0	0	9	1	
<i>Capnocytophaga</i> spp.	3	2	0	1	0	0	0	0	0	0	0	0	0	1	2	
<i>Neisseria</i> spp.	45	35	0	8	2	0	0	0	0	0	0	0	0	38	7	0.279
<i>Fusobacterium</i> spp.	2	1	0	0	2	0	0	0	0	0	0	0	0	2	0	
<i>Borrelia</i> spp.	38	19	0	19	0	0	0	0	1	0	10	26.3	7.5	27	11	0.507
<i>Enterococcus</i> spp.	5	3	0	2	1	0	0	0	1	0	0	0	0	2	3	
<i>Micrococcus</i> spp.	1	1	0	0	0	0	0	0	0	0	0	0	0	1	0	
<i>Gemella</i> spp.	1	0	0	0	1	0	0	0	0	0	0	0	0	1	0	
<i>Peptococcus</i> spp.	1	0	0	1	0	0	1	0	0	0	0	0	0	1	0	
<i>Peptostreptococcus</i> spp.	3	0	0	1	1	1	1	0	0	0	0	0	0	3	0	
<i>Lactobacillus</i> spp.	1	0	0	0	1	0	0	0	0	0	0	0	0	1	0	
Corynebacteria	1	1	0	0	0	0	0	0	0	0	0	0	0	1	0	
<i>Actinomyces</i> spp.	5	0	0	0	4	0	1	0	0	0	0	0	0	4	1	
<i>Escherichia coli</i>	9	6	0	0	0	0	3	0	0	0	0	0	0	1	8	
<i>Enterobacter</i> spp.	3	2	0	1	0	0	0	0	0	0	0	0	0	2	1	
<i>Proteus</i> spp.	2	1	0	0	1	0	0	0	0	0	0	0	0	2	0	
<i>Morganella</i> spp.	3	0	0	1	0	0	2	0	0	0	0	0	0	1	2	
<i>Campylobacter</i> spp.	1	0	0	0	0	0	1	0	0	0	0	0	0	0	1	
<i>Aggregatibacter</i> spp.	3	0	0	0	3	0	0	0	0	0	0	0	0	1	2	
<i>Pseudomonas</i> spp.	7	3	0	1	2	0	0	0	1	0	0	0	0	4	3	
<i>Acinetobacter</i> spp.	4	1	0	3	0	0	0	0	0	0	0	0	0	4	0	
<i>Moraxella</i> spp.	2	0	0	1	0	0	1	0	0	0	0	0	0	2	0	
<i>Sphingomonas</i> spp.	2	2	0	0	0	0	0	0	0	0	0	0	0	2	0	
<i>Prevotella</i> spp.	2	0	0	0	2	0	0	0	0	0	0	0	0	2	0	
<i>Bacteroides</i> spp.	1	0	0	0	0	0	1	0	0	0	0	0	0	1	1	
<i>Xanthomonas</i> spp.	1	0	0	1	0	0	0	0	0	0	0	0	0	1	0	
<i>Aerococcus</i> spp.	1	0	0	0	1	0	0	0	0	0	0	0	0	1	0	
<i>Cryptococcus</i> spp.	27	23	0	4	0	0	0	0	0	0	3	11.1	2.2	21	6	0.999
<i>Candida</i> spp.	5	4	0	1	0	0	0	0	0	0	1	20	0.7	5	0	

Table 2 (continued)

	No.	Meningitis	Encephalitis	Meningoencephalitis	Brain abscess	Subdural empyema	Epidural abscess	Other abscesses	Other syndromes	Arachnoiditis	Chronic infections		<i>p</i> -Value
											Agents ( <i>n</i> , %)	Patients* (%)	
<i>Saccharomyces cerevisiae</i>	1	0	0	0	0	1	0	0	0	0	0	1	0
HSV-1	48	7	21	19	0	0	0	0	0	0	1	2.1	0.060
HSV-2	34	17	6	10	0	0	0	0	1	0	1	2.9	1.000
HSV-1/2	1	1	0	0	0	0	0	0	0	0	0	0	1
EBV	16	9	3	3	10	0	0	0	0	0	3	18.8	0.772
CMV	5	3	0	0	2	0	0	0	0	0	1	20	0.7
VZV	91	17	11	61	0	0	0	0	2	0	0	0	0.000
SFTV	1	0	0	1	0	0	0	0	0	0	0	0	1
WNV	36	19	5	11	10	0	0	0	0	0	3	8.3	0.999
Parechovirus	1	0	1	0	0	0	0	0	0	0	0	0	1
Mumps virus	6	1	0	5	0	0	0	0	0	0	0	0	6
Measles virus	1	0	0	1	0	0	0	0	0	0	0	0	1
Adenovirus	4	3	0	0	0	0	0	0	0	0	10	100	0.7
PIV	2	1	0	1	0	0	0	0	0	0	0	0	4
Enterovirus	91	74	3	15	0	0	0	0	0	0	1	1.1	0.000
HHV6	3	2	1	0	0	0	0	0	0	0	0	0	3
HIV	12	2	4	5	10	0	0	0	0	0	6	50	0.693
JCV	5	0	4	1	0	0	0	0	0	0	2	40	1.5
RSV	1	1	0	0	0	0	0	0	0	0	0	0	1
TBEV	92	1	4	84	0	0	0	0	3	0	0	0	0
<i>Toxoplasma</i> spp.	8	0	0	1	7	0	0	0	0	0	4	50	3
Specified	1079	546	64	388	48	7	23	2	15	1	96	8.9	73
Unspecified	1504	746	144	500	49	7	8	0	54	1	36	2.4	27
Total patients	2583	1292	208	888	97	14	31	2	69	2	132	5.1	0

*Co-NS* Coagulase-negative staphylococci; *HSV* herpes simplex virus; *EBV* Epstein–Barr virus; *CMV* cytomegalovirus; *SFTV* sandfly fever Toscana virus; *WNV* West Nile virus; *PIV* paramyxovirus; *HHV* human herpesvirus; *HIV* human immunodeficiency virus; *JCV* JC virus; *RSV* respiratory syncytial virus; *TBEV* tick-borne encephalitis virus

\*Percentage for chronic CNS disease



**Table 3** The distribution of infecting pathogens according to the ultimate diagnosis

	FR	HU	RO	CZ	DK	HR	BA	KS	SB	AL	IT	PT	TR	SY	IR	KZ	PK	HK	EG	TX	Total	%	Culture(+)		
																							No.	%	
Total	194	188	146	301	192	66	162	87	50	81	53	60	302	80	74	124	114	75	165	69	2583	100	458	100	
<i>S. pneumoniae</i>	25	7	29	6	26	3	4	1	21	2	11	8	34	0	1	14	0	0	0	13	205	8	178	38.9	
<i>S. anginosus</i>	0	0	0	4	10	0	0	0	0	0	0	2	0	0	0	0	4	0	0	0	20	2	17	9.3	
<i>S. sanguinis</i>	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1		1		
<i>S. bovis</i>	1	0	1	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	4		4		
<i>S. mitis</i>	2	1	0	0	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8		4		
<i>S. salivarius</i>	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	2		1		
<i>S. pyogenes</i>	1	2	0	0	0	0	0	0	2	0	1	0	0	0	0	0	0	1	0	0	7		7		
<i>S. viridans</i>	0	1	0	0	1	0	0	1	0	1	0	0	0	0	0	0	0	0	0	1	5		5		
Untyped	0	0	0	1	3	0	0	0	0	0	0	1	2	0	0	0	0	0	0	0	8		4		
<i>S. aureus</i>	1	7	3	5	5	0	2	0	1	1	5	2	4	0	2	1	1	5	0	3	48	1.9	48	10.5	
Co-NS	0	0	1	2	3	0	0	2	1	3	2	3	1	0	1	1	0	3	0	0	23	0.9	23	5.0	
<i>Enterococcus</i> spp.	1	0	2	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	5	*	5	1.1	
<i>Micrococcus luteus</i>	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	*	1	*	
<i>Gemella</i> spp.	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	*	1	*	
<i>Peptococcus</i> spp.	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	*	1	*	
<i>Peptostreptococci</i> spp.	0	0	0	2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	*	3	*	
<i>L. monocytogenes</i>	3	5	0	0	2	1	2	0	2	0	1	3	6	0	0	0	0	2	0	0	27	1	25	5.4	
<i>Lactobacillus</i> spp.	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	*	1	*	
Corynebacteria	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	*	1	*	
<i>Actinomyces</i> spp.	0	0	1	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	4	*	4	*	
<i>Nocardia</i> spp.	1	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	4	*	3	*	
<i>E. coli</i>	0	0	1	1	0	0	0	1	0	0	0	2	2	0	0	0	1	0	0	1	9	1	9	5.4	
<i>Klebsiella</i> spp.	1	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	4	0	0	7		7		
<i>Enterobacter</i> spp.	1	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	3		3		
<i>P. mirabilis</i>	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	2		2		
<i>M. morgani</i>	0	1	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	3		3		
<i>Campylobacter</i> spp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1		1		
<i>H. influenzae</i>	0	0	1	0	0	1	0	0	0	0	3	1	1	0	0	0	0	0	0	0	7	*	7	1.5	
<i>Haemophilus</i> spp.	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	2	*	2	*	
<i>Brucella</i> spp.	0	0	0	0	0	0	0	0	0	2	0	0	7	2	2	0	0	0	0	0	13	0.5	3	*	
<i>Capnocytophaga</i> spp.	2	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	*	2	*	
<i>Aggregatibacter</i> spp.	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	3	*	3	*	
<i>N. meningitidis</i>	3	17	0	2	8	2	1	0	0	1	0	1	6	0	0	0	0	0	1	0	42	1.6	26	5.6	
<i>Kingella kingae</i>	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	*	1	*	
<i>Eikenella corrodens</i>	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	*	2	*	
<i>Pseudomonas</i> spp.	0	1	0	0	1	0	1	0	1	0	0	2	0	0	0	0	1	0	0	0	7	0.6	7	3.3	
<i>Acinetobacter</i> spp.	0	1	0	0	0	0	1	0	1	1	0	0	0	0	0	0	0	0	0	0	4		4		
<i>Moraxella</i> spp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	2		2		
<i>Sphingomonas</i> spp.	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2		2		
<i>Prevotella</i> spp.	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	*	2	*	
<i>B. fragilis</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	*	1	*	
<i>Xanthomonas</i> spp.	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	*	0	0	
<i>Aerococcus urinae</i>	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	*	0	0	
<i>Fusobacterium</i> spp.	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	*	1	*	
<i>M. tuberculosis</i>	3	2	38	0	3	0	14	17	1	6	0	3	25	9	2	0	21	5	0	3	152	5.8	5	1.1	
<i>Cryptococcus</i> spp.	4	0	4	0	0	0	0	0	0	0	4	6	1	0	1	0	0	5	0	2	27	1	20	4.3	
<i>Candida</i> spp.	0	1	0	1	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	0	5	*	4	*	
<i>Saccharomyces cerevisiae</i>	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	*	0	0	
HSV-1	12	3	0	4	9	0	0	0	0	0	4	0	7	3	1	0	1	1	0	3	48		1.8		
HSV-2	10	2	0	4	12	0	0	0	0	0	0	2	1	1	1	0	0	0	0	1	34		1.3		
HSV-1/2	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	*			
EBV	4	2	0	1	1	0	0	0	0	0	3	0	4	0	0	0	0	0	0	1	16		0.6		
CMV	0	4	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	5	*			
VZV	13	1	0	54	11	3	0	0	0	0	2	1	2	0	0	0	0	2	0	2	91		3.5		
SFTV	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	*			
WNV	0	5	0	0	0	2	1	7	0	0	0	0	0	0	0	0	0	18	0	3	36		1.4		
Parechovirus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	*			

**Table 3** (continued)

	FR	HU	RO	CZ	DK	HR	BA	KS	SB	AL	IT	PT	TR	SY	IR	KZ	PK	HK	EG	TX	Total	%	Culture(+)	
																							No.	%
Mumps virus	0	1	0	4	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6	*	
Measles virus	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	*	
Adenovirus	0	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	4	*		
PIV	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	*		
Enterovirus	20	4	0	12	22	8	0	0	0	0	1	8	16	0	0	0	0	0	0	0	91	3.5		
HHV-6	2	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	3	*		
HIV	8	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	9	0.5		
JCV	4	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	5	*		
RSV	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	*		
TBEV	0	14	0	75	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	92	3.5		
<i>Toxoplasma</i> spp.	4	0	0	0	0	0	0	0	0	0	4	0	0	0	0	0	0	0	0	0	8	*		

*Co-NS* Coagulase-negative staphylococci; *SFTV* sandfly fever Toscana virus; *WNV* West Nile virus; *TBEV* tick-borne encephalitis virus; *PIV* parainfluenza virus; *FR* France; *HU* Hungary; *RO* Romania; *CZ* Czech Republic; *DK* Denmark; *HR* Croatia; *BA* Bosnia and Herzegovina; *KS* Kosovo; *SB* Serbia; *AL* Albania; *IT* Italy; *PT* Portugal; *TR* Turkey; *SY* Syria; *IR* Iran; *KZ* Kazakhstan; *PK* Pakistan; *HK* Hong Kong; *EG* Egypt; *TX* Texas

\*The percentage was not specified for <0.005

TBEV [13/250–79/829 ( $p = 0.032$ )] affected adults significantly more than the elderly (Table 2).

### Chronicity

96 (8.9%) out of 1079 patients with any identified pathogen presented with clinical features of chronic meningitis. *Treponema pallidum* (24/96–0/983;  $p < 0.001$ ), *Brucella* spp. (6/96–8/983;  $p = 0.001$ ), *M. tuberculosis* (37/96–115/983;  $p < 0.001$ ), and *Borrelia* spp. (10/96–28/983;  $p = 0.001$ ) were more likely to be causing chronic infections, while streptococci other than pneumococci (0/96–55/983;  $p = 0.012$ ), *Neisseria meningitidis* (0/96; 45/983;  $p = 0.046$ ), VZV (0/96–91/983;  $p = 0.003$ ), enteroviruses (1/96–90/983;  $p = 0.011$ ), and TBEV (0/96–92/983;  $p < 0.001$ ) frequently presented non-chronic patterns. *Streptococcus pneumoniae* (0/96–206/983;  $p < 0.001$ ) always presented with acute patterns (Table 2).

### Immunosuppressive states

There were 741 immunosuppressive conditions in 603 (23.3%) out of 2583 patients. Of these, there were 170 (6.6%) confirmed (primary and secondary) immunosuppressive patients and 571 (22.1%) probable immunosuppressive conditions in 508 cases. The distribution was as follows:

1. Primary immunosuppression: One hyper IgM syndrome (type 1), two hypogammaglobulinemia, one mucocutaneous candidiasis, one IgA deficiency.
2. Secondary immunosuppression: Acute lymphocytic leukemia, bone marrow transplantation, amyloidosis, acute myelocytic leukemia, autoimmune hemolytic anemia, leukemia, myelofibrosis, pyoderma gangrenosum, multiple myelomas, and myelodysplastic syndrome were seen

in one patient. Two monoclonal gammopathies, two chronic lymphocytic leukemias, two large cell lymphomas, three cirrhosis, six solid organ transplantations, seven splenectomies, 31 with chemotherapy, and 121 systemic corticosteroid use were other conditions.

3. Probable immunosuppression: 263 diabetes mellitus, 116 malignancies, 84 HIV infections, 27 pregnancies, 15 alcoholics, ten chronic liver diseases, six Addison's disease, six end-stage renal disease, seven systemic lupus erythematosus, three rheumatoid arthritis, three intravenous drug users, two ulcerative colitis, and two psoriasis. Hypersplenism, acute renal failure, Behçet's disease, positive anti-GM/GSF antibody, brain tumor, congenital leukodystrophy, disseminated sclerosis, Down syndrome, generalized herpetic infection, genital herpes, Crohn's disease, non-Hodgkin lymphoma, panhypopituitarism, scleroderma, and ankylosing spondylitis were seen in one patient.

When the patients with a etiologically proven agent was considered ( $n = 1179$ ), *S. aureus* was more frequent in diabetics (11/121, 9.1%) compared to non-diabetics (37/958, 3.9%) ( $p = 0.017$ ). Coagulase-negative staphylococci were more common in patients with secondary immunosuppression (6/95, 6.3%) than the others (18/984, 1.8%) ( $p = 0.015$ ) and in patients with concurrent malignancies ( $n = 4/61$ , 6.6%) compared to those without concurrent malignancies ( $n = 20/1018$ , 2%) ( $p = 0.042$ ). *Listeria monocytogenes* was more common in secondary immunosuppression (9/95, 9.5%) than those without (18/984, 1.8%) ( $p = 0.015$ ) and in malignancy patients ( $n = 5/61$ , 8.2%) compared to those without malignancy (22/1018, 2.2%) ( $p = 0.015$ ). Finally, cryptococci was the most frequent in etiologically proven CNS infection patients with HIV infection ( $n = 13/50$ , 26%) compared to HIV-negative patients ( $n = 14/1029$ , 1.4%) ( $p < 0.001$ ).

**Table 4** The distribution of ultimate microbiological diagnoses according to the immunosuppression status of the patients

	No.	Probable immunosuppressive conditions					Subtotal	Confirmed immunosuppressive conditions		
		Mal	DM	HIV	Pg	Other		Primary	Secondary	Subtotal
<i>T. pallidum</i>	24	0	0	0	0	0	0	0	0	0
<i>Brucella</i> spp.	14	1	0	0	0	0	1	0	0	0
<i>M. tuberculosis</i>	152	6	18	9	2	6	41	1	10	11
<i>S. pneumoniae</i>	206	14	27	3	3	7	54	1	25	26
<i>Streptococcus</i> spp. (other)	55	1	6	1	1	6	15	0	3	3
<i>S. aureus</i>	48	2	11	0	0	4	17	0	5	5
Co-NS	23	4	4	0	0	1	5	0	6	6
<i>Listeria</i> spp.	27	5	4	0	0	3	12	0	9	9
<i>Nocardia</i> spp.	4	0	0	0	0	0	0	0	2	2
<i>Klebsiella</i> spp.	7	2	1	0	0	0	3	0	0	0
<i>Haemophilus</i> spp.	10	0	1	0	0	0	1	0	1	1
<i>Capnocytophaga</i> spp.	3	1	0	0	0	0	1	0	1	1
<i>Neisseria</i> spp. <sup>a</sup>	45	1	3	0	1	0	5	0	4	4
<i>Fusobacterium</i> spp.	2	0	0	0	0	0	0	0	0	0
<i>Borrelia</i> spp.	38	2	3	0	0	0	5	0	2	2
<i>Enterococcus</i> spp.	5	0	0	0	0	0	0	0	1	1
<i>Micrococcus</i> spp.	1	0	0	0	0	0	0	0	0	0
<i>Gemella</i> spp.	1	0	0	0	0	1	1	0	0	0
<i>Peptococcus</i> spp.	1	0	0	0	0	0	0	0	0	0
<i>Peptostreptococcus</i> spp.	3	0	1	0	0	0	1	0	0	0
<i>Lactobacillus</i> spp.	1	0	1	0	0	0	1	0	0	0
Corynebacteria	1	1	0	0	0	0	1	0	0	0
<i>Actinomyces</i> spp.	5	0	2	0	0	0	2	0	0	0
<i>Escherichia coli</i>	9	2	3	0	0	0	5	0	0	0
<i>Enterobacter</i> spp.	3	1	0	0	0	0	1	0	0	0
<i>Proteus</i> spp.	2	0	0	0	0	0	0	0	0	0
<i>Morganella</i> spp.	3	0	2	0	0	0	2	0	0	0
<i>Campylobacter</i> spp.	1	0	1	0	0	0	1	0	0	0
<i>Aggregatibacter</i> spp.	3	0	0	0	0	0	0	0	0	0
<i>Pseudomonas</i> spp.	7	0	1	0	0	0	1	0	0	0
<i>Acinetobacter</i> spp.	4	1	1	0	0	0	2	0	0	0
<i>Moraxella</i> spp.	2	1	0	0	0	0	1	0	0	0
<i>Sphingomonas</i> spp.	2	2	0	0	0	0	2	0	4	4
<i>Prevotella</i> spp.	2	0	1	0	0	0	1	0	0	0
<i>Bacteroides</i> spp.	1	0	0	0	0	0	0	0	1	1
<i>Xanthomonas</i> spp.	1	0	0	0	0	0	0	0	1	1
<i>Aerococcus</i> urinae	1	0	0	0	0	0	0	0	0	0
<i>Cryptococcus</i> spp.	27	2	1	13	0	1	17	1	5	6
<i>Candida</i> spp.	5	0	2	0	0	1	3	1	0	1
<i>Saccharomyces cerevisiae</i>	1	0	0	0	0	0	0	0	0	0
HSV-1	48	1	6	1	1	1	9	0	3	3
HSV-2	34	3	3	0	0	0	6	0	5	5
HSV-1/2	1	0	0	0	0	0	0	0	0	0
EBV	16	2	3	3	0	0	8	0	3	3
CMV	5	0	0	1	0	0	1	0	1	1
VZV	91	7	12	1	2	2	24	1	10	11
SFTV	1	0	0	0	0	0	0	0	0	0

**Table 4** (continued)

	No.	Probable immunosuppressive conditions						Confirmed immunosuppressive conditions		
		Mal	DM	HIV	Pg	Other	Subtotal	Primary	Secondary	Subtotal
WNV	36	0	7	1	0	0	8	0	0	0
Parechovirus	1	0	0	0	0	0	0	0	0	0
Mumps virus	6	0	0	0	0	0	0	0	0	0
Measles virus	1	0	0	0	0	0	0	0	0	0
Adenovirus	4	0	0	0	0	0	0	0	0	0
PIV	2	0	0	1	0	0	1	0	0	0
Enterovirus	91	1	4	0	0	1	6	0	3	3
HHV-6	3	1	0	0	0	0	1	0	1	1
HIV	12	0	1	12	0	0	13	0	0	0
JCV	5	0	0	5	0	0	5	0	0	0
RSV	1	0	0	1	0	0	1	0	0	0
TBEV	92	1	11	0	0	2	14	0	3	3
<i>Toxoplasma</i> spp.	8	0	1	8	0	0	9	0	0	0
Unspecified	1504	55	142	34	17	15	263	1	70	71
Total patients	2583	116	263	84	27	47	508	5	165	170

*Co-NS* Coagulase-negative staphylococci; *SFTV* sandfly fever Toscana virus; *WNV* West Nile virus; *TBEV* tick-borne encephalitis virus; *PIV* parainfluenza virus; *Mal* malignancy; *DM* diabetes mellitus; *Pg* pregnancy; *Primary* and *secondary* indicate primary and secondary immunodeficiencies, respectively


<sup>a</sup> *Eikenella* and *Kingella* were included in this group

**Outcomes**

In this study, 477 (18.5%) patients survived with sequelae and 227 (8.8%) died during hospital stay. Motor deficit (*n* = 128, 5%), cranial nerve involvement (*n* = 92, 3.6%), cognitive impairment (*n* = 70, 2.7%), persisting headache (*n* = 38, 1.5%), and epilepsy (*n* = 26, 1%) were the most frequent sequelae. Overall, 1879 (72.7%) patients were discharged with complete cure. Among the patients with any etiologically proven pathogen, 711 (65.9%) patients

had complete cure, 261 (24.2%) survived with sequelae, and 107 (9.9%) patients died. Adverse clinical outcomes in patients with an etiologically proven pathogen [368/1079(34.1%)] were significantly more common than those without confirmed etiological diagnosis [336/1504 (22.7%)] (*p* < 0.001). The Glasgow Coma Scale (GCS) scores of the patients with a etiologically proven CNS pathogen on admission were significantly lower in patients with adverse clinical outcomes (mean: 11.94 ± 3.69) compared to survivors with complete cure (mean: 13.37 ± 2.86)

**Fig. 1** Common pathogens in community-acquired central nervous system (CA-CNS) infections



( $p < 0.001$ ). The outcome distributions in CA-CNS infection patients are presented in Table 5.

- a) Mortality rate by specific pathogens: The fatality rate was significantly higher for *L. monocytogenes* [(8/107–19/972); ( $p = 0.003$ )], *S. aureus* [(10/107–38/972); ( $p = 0.021$ )], *M. tuberculosis* [(28/107–124/972); ( $p < 0.001$ )], and *S. pneumoniae* [(33/107–173/972); ( $p = 0.002$ )], while it was significantly lower in enterovirus [(1/107–90/972); ( $p = 0.006$ )] and TBEV [(1/107–91/972); ( $p = 0.005$ )] infections, and where no death was seen due to *Borrelia* spp. infections [(0/107–38/972); (non-applicable statistics)].
- b) Rate of neurological sequelae by specific pathogens: Persistence of sequelae was significantly higher in herpes simplex virus (HSV) type 1 cases [(24/261–24/818); ( $p < 0.001$ )], while it was less frequent in enterovirus [(13/261–78/818); ( $p = 0.029$ )] and TBEV [(8/261–84/818); ( $p < 0.001$ )] infections.

## Discussion

CNS infections have significant mortality and permanent neurological sequelae worldwide [17]. Despite the advances in management, 27.3% of patients with CA-CNS infections experienced unfavorable outcomes in this study, including persistent sequelae and death. The most frequent sequelae were motor deficits (5%) and cranial nerve involvements (3.6%), in accordance with the current literature [19]. Cognitive impairment, persisting headache, and epilepsy were seen in descending order (Table 5). We found that CA-CNS diseases due to *S. aureus*, *M. tuberculosis*, *S. pneumoniae*, and *Listeria* spp. were the most fatal etiologies, while the patients with CA-CNS infections due to enterovirus, TBEV, and *Borrelia* spp. were less likely to die. Furthermore, HSV-1 was associated with sequelae most commonly, while enteroviruses and TBEV were less likely to be complicated with sequelae.

Meningitis and encephalitis, known as two intersecting syndromes, were reported to be the most frequent CNS infections [17, 20, 21]. The use of neuroimaging before lumbar puncture has generated considerable debate, with some investigators pointing to delays in antibiotic administration, reduced likelihood of identifying a pathogen, and an increase in mortality [22]. In this study, half of the cases were classified as meningitis and 8% as encephalitis, and an unspecified 34% of patients were classified as meningoencephalitis. Considering the high frequency of cases presenting with encephalitic pattern in this study, the detection of brain parenchymal involvement with imaging together with positive EEG

findings is directly suggestive of encephalitis [23] and leads to early administration of antivirals that reduce unfavorable outcomes [8]. Added to that, suppurative intracranial infections, where brain abscesses dominated in two-thirds and epidural abscesses in one-third, made up 5.6% of all CA-CNS infections. Thus, these data indicate the necessity of using radioimaging methods and EEG in an unspecified CA-CNS infection patient considering the high frequency encephalitis and suppurative intracranial infections.

Consequently, the diagnostic capacity of the hospitals treating CA-CNS infections determines the early diagnosis, which is key to rational therapies [2, 4, 8, 17]. We found that the etiological diagnosis was established in 41.8% of all patients. The data are scarce in the literature in that context. In an African study, 35% of the cases with meningitis [24] and in a Turkish systematic review, 32.6% of the cases [25] were reported to have an etiological diagnosis. In addition, 3% of the CA-CNS infections had multiple infecting agents, which must not be overlooked by the treating clinicians. The most frequent infecting pathogens in this study were *S. pneumoniae* and *M. tuberculosis*. TBEV, VZV, enteroviruses, and HSV infections are among the other most frequent CA-CNS infections. Hence, these data highlight the importance of pneumococcal immunization, producing herd immunity and decreasing invasive disease [22], and widespread tuberculosis elimination programs. Accordingly, *N. meningitidis*, which is the agent of epidemic meningitis, was a still noteworthy pathogen and had a share of 1.6% in CA-CNS patients. Although *Streptococcus suis* was reported to be one of the major causes of meningitis in some parts of Asia [22], *Streptococcus anginosus* was recovered mainly from patients in European countries among the non-pneumococcal species in this study. In a recent meta-analysis, the most frequent causes of brain abscesses were streptococci and *S. aureus* [26], and our data were in accordance with this. Pneumococcal CNS disease and VZV infections were the most frequent infections in the elderly, followed by *M. tuberculosis* and *L. monocytogenes* infections. In the literature, pneumococcal and listerial CNS diseases have been known to be quite frequent in elderly patients [4, 27]. However, our data affirm that, in accordance with the high tuberculous CNS disease worldwide [28], and as specific cell-mediated immunity declines in the elderly for VZV infections [29], they should not be underestimated in the elderly. According to our data, *S. pneumoniae*, *S. aureus*, *L. monocytogenes*, and VZV infections were seen relatively more frequently in older patients, while *M. tuberculosis* and enterovirus infections were observed in adult CA-CNS cases. These differences may be explained owing to the decreasing immunity with advancing age and epidemiological trends [4, 21, 30].

*Listeria monocytogenes* meningitis was more often found in patients with acquired immunodeficiencies [2, 31], while cryptococcal CNS disease was common in HIV-infected

**Table 5** The outcomes of the CNS infection patients

	Cure		Cure with sequelae									Death	GCS <sup>a</sup>	
	No.	n, %	CNI	Epilepsy	Ataxia	MD	UI	CI	HA	Other <sup>b</sup>	n (%)	Mean	SD	
<b>Bacterial CNS infections</b>														
<i>T. pallidum</i>	24	14	10 (42)	0	1	0	1	0	0	0	8	0	14.54	1.14
<i>Brucella</i> spp.	14	12	2 (14)	0	0	1	1	0	0	0	0	0	12.25	3.16
<i>M. tuberculosis</i>	152	90	34 (22)	9	3	1	10	0	0	1	10	28 (18.4)	12.56	3.18
<i>S. pneumoniae</i>	206	128	45 (22)	9	0	1	10	2	5	2	16	33 (16)	10.31	3.57
<i>Streptococcus</i> spp. (other)	55	31	17 (31)	2	2	0	8	0	3	1	1	7 (13)	12.44	3.1
<i>S. aureus</i>	48	24	14 (29)	1	2	0	5	0	1	0	5	10 (21)	12.72	3.22
Staphylococci (other)	23	19	4 (17)	1	0	0	1	1	0	0	1	0	12.27	3.85
<i>Listeria</i> spp.	27	15	4 (15)	0	0	0	1	0	2	0	1	8 (30)	10.65	4.01
<i>Nocardia</i> spp.	4	2	1 (25)	0	0	0	0	1	0	0	0	1 (25)	15	0
<i>Klebsiella</i> spp.	7	5	0	0	0	0	0	0	0	0	0	2 (29)	12.71	3.94
<i>Haemophilus</i> spp.	10	8	1 (10)	0	0	0	0	0	1	0	0	1 (10)	13.5	2.71
<i>Capnocytophaga</i> spp.	3	2	1 (33)	0	0	0	1	0	0	0	0	0	13	3.46
<i>Neisseria</i> spp. <sup>c</sup>	45	32	8 (18)	0	0	0	0	0	2	4	2	5 (11)	11.98	3.99
<i>Fusobacterium</i> spp.	2	1	1 (50)	0	0	0	1	0	0	0	0	0	14.5	0.70
<i>Borrelia</i> spp.	38	25	13 (34)	6	0	0	1	0	0	0	6	0	14.27	2.14
<i>Enterococcus</i> spp.	5	3	2 (40)	0	0	0	1	0	1	0	0	0	11	1.41
<i>Micrococcus</i> spp.	1	1	0	0	0	0	0	0	0	0	0	0	14	0
<i>Gemella</i> spp.	1	0	1 (100)	0	1	0	0	0	0	0	0	0	15	0
<i>Peptococcus</i> spp.	1	1	0	0	0	0	0	0	0	0	0	0	8	0
<i>Peptostreptococcus</i> spp.	3	1	2 (67)	0	0	0	2	0	0	0	0	0	10.33	4.04
<i>Lactobacillus</i>	1	0	1 (100)	0	0	0	1	0	0	0	0	0	15	0
Corynebacteria	1	0	0	0	0	0	0	0	0	0	0	1 (100)	14	0
<b>Actinomyces</b>														
<i>Escherichia coli</i>	9	6	1 (11)	0	0	0	0	0	0	0	1	2 (22)	12.57	2.99
<i>Enterobacter</i> spp.	3	2	1 (33)	0	0	0	1	0	0	0	0	0	9	8.48
<i>Proteus</i> spp.	2	1	1 (50)	1	0	0	0	0	0	0	0	0	13.5	2.12
<i>Morganella</i> spp.	3	3	0	0	0	0	0	0	0	0	0	0	14.67	0.57
<i>Campylobacter</i> spp.	1	1	0	0	0	0	0	0	0	0	0	0	15	0
<i>Aggregatibacter</i> spp.	3	2	1 (33)	0	0	0	1	0	0	0	0	0	14.67	0.57
<i>Pseudomonas</i> spp.	7	4	2 (29)	0	0	0	0	0	0	2	0	1 (14)	12.14	4.56
<i>Acinetobacter</i> spp.	4	2	2 (50)	1	1	0	0	0	0	0	0	0	10.67	4.04
<i>Moraxella</i> spp.	2	1	1 (50)	0	0	0	1	0	0	0	0	0	12	6.36
<i>Sphingomonas</i> spp.	2	2	0	0	0	0	0	0	0	0	0	0	15	0
<i>Prevotella</i> spp.	2	0	2 (100)	0	0	0	2	0	0	0	0	0	15	0
<i>Bacteroides</i> spp.	1	0	1 (100)	0	0	0	1	0	0	0	0	0	13	0
<i>Xanthomonas</i> spp.	1	1	0	0	0	0	0	0	0	0	0	0	15	0
<i>Aerococcus urinae</i>	1	1	0	0	0	0	0	0	0	0	0	0	14	0
<b>Fungal CNS infections</b>														
<i>Saccharomyces cerevisiae</i>	1		1 (100)	0	0	0	1	0	0	0	0	0	10	0
<i>Cryptococcus</i> spp.	27	18	5 (19)	2	1	0	0	0	1	0	1	4 (15)	13.97	2.55
<i>Candida</i> spp.	5	3	0	0	0	0	0	0	0	0	0	2 (40)	11.5	4.04
<b>Viral CNS infections</b>														
HSV-1	48	22	24 (50)	2	3	0	0	0	19	0	0	2 (4)	12.81	3.33
HSV-2	34	25	7 (21)	0	1	0	0	0	4	1	1	2 (6)	14.06	2.13
HSV-1/2	1		1 (100)	0	0	0	0	0	0	0	1	0		
EBV	16	11	4 (25)	1	1	0	1	1	0	0	0	1 (6)	12.75	3.45



**Table 5** (continued)

	Cure		Cure with sequelae									Death	GCS <sup>a</sup>	
	No.	n, %	CNI	Epilepsy	Ataxia	MD	UI	CI	HA	Other <sup>b</sup>	n (%)	Mean	SD	
CMV	5	2	3 (60)	0	0	0	0	1	0	1	1	0	15	0
VZV	91	61	25 (27)	14	0	0	4	0	4	0	3	5 (5.5)	14.39	1.63
SFTV	1	1	0	0	0	0	0	0	0	0	0	0	15	0
WNV	36	24	9 (25)	1	1	0	4	0	1	2	0	3 (8)	13.17	3.47
Parechovirus	1	1	0	0	0	0	0	0	0	0	0	0	15	0
Mumps virus	6	5	1 (17)	1	0	0	0	0	0	0	0	0	15	0
Measles virus	1	1	0	0	0	0	0	0	0	0	0	0	15	0
Adenovirus	4	2	1 (25)	0	0	0	0	0	0	1	0	0	15	0
PIV	2	1	1 (50)	0	0	0	0	0	0	0	1	0	15	0
Enterovirus	91	77	13 (14)	1	0	0	0	0	3	9	0	1 (1.1)	14.72	1.2
HHV-6	3	3	0	0	0	0	0	0	0	0	0	0	13	3.46
HIV	12	5	7 (58)	2	0	0	0	0	5	0	0	0	13.25	3.59
JCV	5	1	4 (80)	0	0	0	1	0	2	0	1	0	14.2	1.30
RSV	1	1	0	0	0	0	0	0	0	0	0	0	15	0
TBEV	92	83	8 (8.6)	3	0	0	3	0	1	0	1	1 (1.1)	14.57	0.95
<i>Toxoplasma</i> spp.	8	5	3 (38)	0	0	0	0	0	1	0	2	0	11.88	4.96
Undermined etiology	1504	1168	216 (14)	35	9	3	62	1	14	14	78	120 (8)		
Total	2583	1879	477 (18)	92	26	6	128	7	70	38	141	227 (9)		

CNI Cranial nerve involvement; MD motor deficit; UI urinary incontinency; CI cognitive impairment; HA persisting headache; GCS Glasgow Coma Scale; SD standard deviation; SFTV sandfly fever Toscana virus; WNV West Nile virus; TBEV tick-borne encephalitis virus; PIV parainfluenza virus

<sup>a</sup> On admission data

<sup>b</sup> Depression, spinal abscess, cerebellar syndrome, etc.

<sup>c</sup> *Eikenella* and *Kingella* were included in this group

patients [32, 33]. Our data were in accordance with that for both pathogens. In addition, we have disclosed that *S. aureus* was a significant agent in diabetic patients, while coagulase-negative staphylococci were common in patients with secondary immunosuppression and malignancy patients. These two groups of pathogens were reported to be the leading agents in various subgroups of immunocompromised patients [34, 35]. However, to the best of our knowledge, this is the first report interrelating immunosuppressive states and staphylococcal CA-CNS infections. Our data may, therefore, support the recommendations in using anti-Gram-positive agents as part of the combination regimen in bacterial CA-CNS infections [2, 3, 7].

In the literature of the past two decades [32, 33, 36], the most frequent agents causing chronic CNS infections were cryptococci and *M. tuberculosis*, depending on the HIV status. In HIV-positive patients, cryptococcal disease predominated [32, 33] and in HIV-negative patients, CNS tuberculosis [36] was the most frequent diagnosis in chronic CNS infections. In this study, 4.3% of the cases presented with a chronic pattern extending over 4 weeks, and neurosyphilis, brucellar meningitis, neuroborreliosis, and CNS tuberculosis were significantly more likely to present chronic courses. In contrast,

cryptococcal meningitis was seen only in 2.2% of cases with chronic CNS disease, although it was still the most frequent CA-CNS infection in HIV-positive individuals. This shift was most probably due to the increasing efficiency of HIV treatment worldwide, with the resultant decrease in opportunistic infections [37].

We believe that our study has important implications for the management of CA-CNS infections. *Streptococcus pneumoniae* and *M. tuberculosis* were the leading CA-CNS pathogens. Considering the high frequency of patients predisposed as encephalitis, meningoencephalitis, or suppurative intracranial infections, both MRI and EEG should be applied to all patients with probable CA-CNS infections. The need to improve the diagnostic capacity in centers treating CNS infections is another serious concern. Accordingly, high sequelae forming infections like HSV-1 were important in the differential diagnosis. In addition to pneumococcal disease and CNS tuberculosis, both common in all ages, VZV, *S. aureus*, and *L. monocytogenes* should be taken into consideration in the elderly population. Accordingly, staphylococcal, cryptococcal, and listerial infections should be considered for immunocompromised patients. Finally, clinicians need to consider that neurosyphilis, brucellar meningitis,

neuroborreliosis, and CNS tuberculosis may have a predilection for prolonged courses where alternative strategies are needed.

#### Compliance with ethical standards

**Funding** We did not receive any kind of funding.

**Ethical approval** Yes, it is obtained from the Review Board of Fatih Sultan Mehmet Education and Research Hospital in Istanbul.

**Informed consent** Not applicable. The study has a retrospective design.

**Conflict of interest** We have no competing interests to declare.

#### References

- McGill F, Heyderman RS, Michael BD, Defres S, Beeching NJ, Borrow R, Glennie L, Gaillemain O, Wyncoll D, Kaczmarski E, Nadel S, Thwaites G, Cohen J, Davies NW, Miller A, Rhodes A, Read RC, Solomon T (2016) The UK joint specialist societies guideline on the diagnosis and management of acute meningitis and meningococcal sepsis in immunocompetent adults. *J Infect* 72(4):405–438
- van de Beek D, Cabellos C, Dzapova O, Esposito S, Klein M, Kloek AT, Leib SL, Mourvillier B, Ostergaard C, Pagliano P, Pfister HW, Read RC, Sipahi OR, Brouwer MC; ESCMID Study Group for Infections of the Brain (ESGIB) (2016) ESCMID guideline: diagnosis and treatment of acute bacterial meningitis. *Clin Microbiol Infect* 22(Suppl 3):S37–S62
- Tunkel AR, Glaser CA, Bloch KC, Sejvar JJ, Marra CM, Roos KL, Hartman BJ, Kaplan SL, Scheld WM, Whitley RJ; Infectious Diseases Society of America (2008) The management of encephalitis: clinical practice guidelines by the Infectious Diseases Society of America. *Clin Infect Dis* 47(3):303–327
- Tunkel AR, Hartman BJ, Kaplan SL, Kaufman BA, Roos KL, Scheld WM, Whitley RJ (2004) Practice guidelines for the management of bacterial meningitis. *Clin Infect Dis* 39(9):1267–1284
- Solomon T, Michael BD, Smith PE, Sanderson F, Davies NW, Hart JJ, Holland M, Easton A, Buckley C, Kneen R, Beeching NJ; National Encephalitis Guidelines Development and Stakeholder Groups (2012) Management of suspected viral encephalitis in adults—Association of British Neurologists and British Infection Association National Guidelines. *J Inf Secur* 64(4):347–373
- Erdem H, Ozturk-Engin D, Tireli H, Kilicoglu G, Defres S, Gulsun S, Sengoz G, Crisan A, Johansen IS, Inan A, Nechifor M, Al-Mahdawi A, Civljak R, Ozguler M, Savic B, Ceran N, Cacopardo B, Inal AS, Namiduru M, Dayan S, Kayabas U, Parlak E, Khalifa A, Kursun E, Sipahi OR, Yemisen M, Akbulut A, Bitirgen M, Popovic N, Kandemir B, Luca C, Parlak M, Stahl JP, Pehlivanoglu F, Simeon S, Ulu-Kilic A, Yasar K, Yilmaz G, Yilmaz E, Beovic B, Catroux M, Lakatos B, Sunbul M, Oncul O, Alabay S, Sahin-Horasan E, Kose S, Shehata G, Andre K, Dragovac G, Gul HC, Karakas A, Chadapaud S, Hansmann Y, Harxhi A, Kirova V, Masse-Chabredier I, Oncu S, Sener A, Tekin R, Elaldi N, Deveci O, Ozkaya HD, Karabay O, Senbayrak S, Agalar C, Vahaboglu H (2015) Hamsi scoring in the prediction of unfavorable outcomes from tuberculous meningitis: results of Haydarpasa-II study. *J Neurol* 262(4):890–898
- Erdem H, Elaldi N, Öztoprak N, Sengoz G, Ak O, Kaya S, Inan A, Nayman-Alpat S, Ulu-Kilic A, Pekok AU, Gunduz A, Gozel MG, Pehlivanoglu F, Yasar K, Yilmaz H, Hatipoglu M, Cicek-Senturk G, Akcam FZ, Inkaya AC, Kazak E, Sagmak-Tartar A, Tekin R, Ozturk-Engin D, Ersoy Y, Sipahi OR, Guven T, Tuncer-Ertem G, Alabay S, Akbulut A, Balkan II, Oncul O, Cetin B, Dayan S, Ersoz G, Karakas A, Ozgunes N, Sener A, Yesilkaya A, Erturk A, Gundes S, Karabay O, Sirmatel F, Tosun S, Turhan V, Yalci A, Akkoyunlu Y, Aydin E, Diktas H, Kose S, Ulcay A, Seyman D, Savasci U, Leblebicioglu H, Vahaboglu H (2014) Mortality indicators in pneumococcal meningitis: therapeutic implications. *Int J Infect Dis* 19: 13–19
- Erdem H, Cag Y, Ozturk-Engin D, Defres S, Kaya S, Larsen L, Poljak M, Barsic B, Argemi X, Sørensen SM, Bohr AL, Tattevin P, Gunst JD, Baštáková L, Jereb M, Johansen IS, Karabay O, Pekok AU, Sipahi OR, Chehri M, Beraud G, Shehata G, Del Vecchio RF, Maresca M, Karsen H, Sengoz G, Sunbul M, Yilmaz G, Yilmaz H, Sharif-Yakan A, Kanj SS, Parlak E, Pehlivanoglu F, Korkmaz F, Komur S, Kose S, Ulug M, Bolukcu S, Coskuner SA, Ince N, Akkoyunlu Y, Halac G, Sahin-Horasan E, Tireli H, Kilicoglu G, Al-Mahdawi A, Nemli SA, Inan A, Senbayrak S, Stahl JP, Vahaboglu H (2015) Results of a multinational study suggest the need for rapid diagnosis and early antiviral treatment at the onset of herpetic meningoencephalitis. *Antimicrob Agents Chemother* 59(6):3084–3089
- Thwaites G, Fisher M, Hemingway C, Scott G, Solomon T, Innes J; British Infection Society (2009) British Infection Society guidelines for the diagnosis and treatment of tuberculosis of the central nervous system in adults and children. *J Infect* 59(3):167–187
- Helbok R, Broessner G, Pfausler B, Schmutzhard E (2009) Chronic meningitis. *J Neurol* 256(2):168–175
- Zizza CA, Ellison KJ, Wernette CM (2009) Total water intakes of community-living middle-old and oldest-old adults. *J Gerontol A Biol Sci Med Sci* 64(4):481–486
- García JM, Gamboa P, de la Calle A, Hernández MD, Caballero MT, García BE, Labrador M, Lahoz C, Longo Areso N, López Hoyos M, Martínez Quesada J, Mayorga L, Monteseirin FJ, Sanz ML; Committee of Immunology of the Spanish Society of Allergy and Clinical Immunology (2010) Diagnosis and management of immunodeficiencies in adults by allergologists. *J Investig Allergol Clin Immunol* 20(3):185–194
- Centers for Disease Control and Prevention (CDC) (2013) STD Surveillance Case Definitions. Available online at: <http://www.cdc.gov/std/stats/casedefinitions-2014.pdf>
- Halperin JJ, Shapiro ED, Logigian E, Belman AL, Dotevall L, Wormser GP, Krupp L, Gronseth G, Bever CT Jr; Quality Standards Subcommittee of the American Academy of Neurology (2007) Practice parameter: treatment of nervous system Lyme disease (an evidence-based review): report of the quality standards Subcommittee of the American Academy of Neurology. *Neurology* 69(1):91–102
- Erdem H, Kilic S, Sener B, Acikel C, Alp E, Karahocagil M, Yetkin F, Inan A, Kecik-Bosnak V, Gul HC, Tekin-Koruk S, Ceran N, Demirdal T, Yilmaz G, Ulu-Kilic A, Ceylan B, Dogan-Celik A, Nayman-Alpat S, Tekin R, Yalci A, Turhan V, Karaoglan I, Yilmaz H, Mete B, Batirel A, Ulcay A, Dayan S, Seza Inal A, Ahmed SS, Tufan ZK, Karakas A, Teker B, Namiduru M, Savasci U, Pappas G (2013) Diagnosis of chronic brucellar meningitis and meningoencephalitis: the results of the Istanbul-2 study. *Clin Microbiol Infect* 19(2):E80–E86
- Erdem H, Ozturk-Engin D, Elaldi N, Gulsun S, Sengoz G, Crisan A, Johansen IS, Inan A, Nechifor M, Al-Mahdawi A, Civljak R, Ozguler M, Savic B, Ceran N, Cacopardo B, Inal AS, Namiduru M, Dayan S, Kayabas U, Parlak E, Khalifa A, Kursun E, Sipahi OR, Yemisen M, Akbulut A, Bitirgen M, Dulovic O, Kandemir B, Luca C, Parlak M, Stahl JP, Pehlivanoglu F, Simeon S, Ulu-Kilic A,

- Yasar K, Yilmaz G, Yilmaz E, Beovic B, Catroux M, Lakatos B, Sunbul M, Oncul O, Alabay S, Sahin-Horasan E, Kose S, Shehata G, Andre K, Alp A, Cosic G, Cem Gul H, Karakas A, Chadapaud S, Hansmann Y, Harxhi A, Kirova V, Masse-Chabredier I, Oncu S, Sener A, Tekin R, Deveci O, Karabay O, Agalar C (2014) The microbiological diagnosis of tuberculous meningitis: results of Haydarpasa-1 study. *Clin Microbiol Infect* 20(10):O600–O608
17. Tunkel AR (2015) Approach to the patient with central nervous system infection. In: Bennett JE, Dolin R, Blaser MJ (eds) Mandell, Douglas, and Bennett's principles and practice of infectious diseases. Elsevier, Philadelphia, pp 1091–1096
  18. Sánchez-Portocarrero J, Pérez-Cecilia E, Corral O, Romero-Vivas J, Picazo JJ (2000) The central nervous system and infection by *Candida* species. *Diagn Microbiol Infect Dis* 37(3):169–179
  19. Lucas MJ, Brouwer MC, van de Beek D (2016) Neurological sequelae of bacterial meningitis. *J Infect* 73(1):18–27
  20. Beckham JD, Tyler KL (2015) Encephalitis. In: Bennett JE, Dolin R, Blaser MJ (eds) Mandell, Douglas, and Bennett's principles and practice of infectious diseases. Elsevier, Philadelphia, pp 1144–1163
  21. Tunkel AR, van de Beek D, Scheld MW (2015) Acute meningitis. In: Bennett JE, Dolin R, Blaser MJ (eds) Mandell, Douglas, and Bennett's principles and practice of infectious diseases. Elsevier, Philadelphia, pp 1097–1137
  22. McGill F, Heyderman RS, Panagiotou S, Tunkel AR, Solomon T (2016) Acute bacterial meningitis in adults. *Lancet* 388(10063):3036–3047
  23. Cag Y, Erdem H, Leib S, Defres S, Kaya S, Larsen L, Poljak M, Ozturk-Engin D, Barsic B, Argemi X, Sørensen SM, Bohr AL, Tattevin P, Gunst JD, Baštáková L, Jereb M, Johansen IS, Karabay O, Pekok AU, Sipahi OR, Chehri M, Beraud G, Shehata G, Fontana R, Maresca M, Karsen H, Sengoz G, Sunbul M, Yilmaz G, Yilmaz H, Sharif-Yakan A, Kanj S, Parlak E, Pehlivanoglu F, Korkmaz F, Komur S, Kose S, Ulug M, Bolukcu S, Coskuner SA, Stahl JP, Ince N, Akkoyunlu Y, Halac G, Sahin-Horasan E, Tireli H, Kilicoglu G, Al-Mahdawi A, Nemli SA, Inan A, Senbayrak S, Vahaboglu H, Elaldi N (2016) Managing atypical and typical herpetic central nervous system infections: results of a multinational study. *Clin Microbiol Infect* 22(6):568.e9–568.e17
  24. Nuoh RD, Nyarko KM, Nortey P, Sackey SO, Lwanga NC, Ameme DK, Nuolabong C, Abdulai M, Wurapa F, Afari E (2016) Review of meningitis surveillance data, upper West region, Ghana 2009–2013. *Pan Afr Med J* 25(Suppl 1):9. doi:10.11604/pamj.supp.2016.25.1.6180
  25. Arda B, Sipahi OR, Atalay S, Ulusoy S (2008) Pooled analysis of 2,408 cases of acute adult purulent meningitis from Turkey. *Med Princ Pract* 17(1):76–79
  26. Brouwer MC, Coutinho JM, van de Beek D (2014) Clinical characteristics and outcome of brain abscess: systematic review and meta-analysis. *Neurology* 82(9):806–813
  27. Erdem H, Kilic S, Coskun O, Ersoy Y, Cagatay A, Onguru P, Alp S; Members of the Turkish Bacterial Meningitis in the Elderly Study Group (2010) Community-acquired acute bacterial meningitis in the elderly in Turkey. *Clin Microbiol Infect* 16(8):1223–1229
  28. World Health Organization (WHO) (2015) Global tuberculosis report 2015, 20th edition. Available online at: [http://apps.who.int/iris/bitstream/10665/191102/1/9789241565059\\_eng.pdf?ua=1](http://apps.who.int/iris/bitstream/10665/191102/1/9789241565059_eng.pdf?ua=1)
  29. Gilden D, Nagel M, Cohrs R, Mahalingam R, Baird N (2015) Varicella zoster virus in the nervous system. *F1000Res* 4:F1000 Faculty Rev-1356
  30. Skoczyńska A, Kuch A, Sadowy E, Waško I, Markowska M, Ronkiewicz P, Matynia B, Bojarska A, Wasiak K, Gołębiowska A, van der Linden M, Hryniewicz W; Participants of a laboratory-based surveillance of community acquired invasive bacterial infections (BINet) (2015) Recent trends in epidemiology of invasive pneumococcal disease in Poland. *Eur J Clin Microbiol Infect Dis* 34(4):779–787
  31. Koopmans MM, Brouwer MC, Bijlsma MW, Bovenkerk S, Keijzers W, van der Ende A, van de Beek D (2013) *Listeria monocytogenes* sequence type 6 and increased rate of unfavorable outcome in meningitis: epidemiologic cohort study. *Clin Infect Dis* 57(2):247–253
  32. Mihret W, Zenebe G, Bekele A, Abebe M, Wassie L, Yamuah LK, Woldemeskel D, Kassahun Y, Medhin G, Engers H, Aseffa A (2014) Chronic meningitis in immunocompromised adult Ethiopians visiting Tikur Anbessa teaching hospital and Ye'huleshet clinic from 2003–2004. *Ethiop Med J Suppl* 1:43–48
  33. Helbok R, Pongpakdee S, Yenjun S, Dent W, Beer R, Lackner P, Bunyaratvej P, Prasert B, Vejajjiva A, Schmutzhard E (2006) Chronic meningitis in Thailand. Clinical characteristics, laboratory data and outcome in patients with specific reference to tuberculosis and cryptococcosis. *Neuroepidemiology* 26(1):37–44
  34. Taramasso L, Tatarelli P, Di Biagio A (2016) Bloodstream infections in HIV-infected patients. *Virulence* 7(3):320–328
  35. Heidenreich D, Kreil S, Nolte F, Reinwald M, Hofmann WK, Klein SA (2016) Allogeneic hematopoietic cell transplantation without fluconazole and fluoroquinolone prophylaxis. *Ann Hematol* 95(2):287–293
  36. Anderson NE, Willoughby EW (1987) Chronic meningitis without predisposing illness—a review of 83 cases. *Q J Med* 63(240):283–295
  37. Baxter C, Moodley D (2015) Improving adolescent maternal health. *S Afr Med J* 105(11):948–952